


**Universität
Stuttgart**



On the well-posedness of entropy solutions to
conservation laws with a zero-flux boundary condition

Raimund Bürger, Hermano Frid, Kenneth H. Karlsen

**Berichte aus dem Institut für
Angewandte Analysis und Numerische Simulation**

Preprint 2004/018

Universität Stuttgart

On the well-posedness of entropy solutions to
conservation laws with a zero-flux boundary condition

Raimund Bürger, Hermano Frid, Kenneth H. Karlsen

**Berichte aus dem Institut für
Angewandte Analysis und Numerische Simulation**

Preprint 2004/018

Institut für Angewandte Analysis und Numerische Simulation (IANS)
Fakultät Mathematik und Physik
Fachbereich Mathematik
Pfaffenwaldring 57
D-70 569 Stuttgart

E-Mail: ians-preprints@mathematik.uni-stuttgart.de

WWW: <http://preprints.ians.uni-stuttgart.de>

ISSN **1611-4176**

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.
IANS-Logo: Andreas Klimke. \LaTeX -Style: Winfried Geis, Thomas Merkle.

ON THE WELL-POSEDNESS OF ENTROPY SOLUTIONS TO CONSERVATION LAWS WITH A ZERO-FLUX BOUNDARY CONDITION

RAIMUND BÜRGER^A, HERMANO FRID^B, AND KENNETH H. KARLSEN^C

ABSTRACT. We study a zero-flux type initial-boundary value problem for scalar conservation laws with a genuinely nonlinear flux. We suggest a notion of entropy solution for this problem and prove its well-posedness. The asymptotic behavior of entropy solutions is also discussed.

1. INTRODUCTION

In recent years significant advances have been made in the analysis of initial-boundary value problems for multi-dimensional scalar conservation laws of the type

$$\partial_t u + \nabla_{\mathbf{x}} \cdot \mathbf{f}(u) = 0, \quad (\mathbf{x}, t) \in Q_T := \Omega \times (0, T), \quad (1.1)$$

where $\Omega \subseteq \mathbb{R}^N$ is a bounded spatial domain, $T > 0$, and the flux vector \mathbf{f} is a smooth function of the unknown u . Moreover, (1.1) is supplemented with an initial condition

$$u(\mathbf{x}, 0) = u_0(\mathbf{x}), \quad \mathbf{x} \in \Omega. \quad (1.2)$$

It is well known that solutions of nonlinear conservation laws may become discontinuous as time evolves, even for smooth initial data, such that (1.1) has to be understood in the distributional sense. This in turn requires an entropy condition to select the physically relevant discontinuous solution, called the *entropy solution*.

A well studied boundary condition for (1.1), (1.2) is the Dirichlet boundary condition

$$u(\mathbf{x}, t) = \phi(\mathbf{x}) \quad \text{for } (\mathbf{x}, t) \in \partial\Omega \times (0, T), \text{ e.g. } \phi \in L^\infty(\partial\Omega). \quad (1.3)$$

However, the boundary datum (1.3) may not always provide the most natural setting for conservation laws on bounded domains. For example, assume that u is the local density of a continuous phase that assumes values from a finite interval $[0, u_{\max}]$ only, and is associated with a kinematic flow velocity $\mathbf{v}(u)$. Then a bounded domain Ω typically corresponds to a closed container with impermeable rigid walls that induce the zero-flux boundary condition $(u\mathbf{v}(u)) \cdot \mathbf{n} = 0$ on $\partial\Omega$, where \mathbf{n} is the outer normal vector to the boundary $\partial\Omega$ of Ω . This suggests the alternative zero-flux boundary condition

$$\mathbf{f}(u) \cdot \mathbf{n} = 0 \quad \text{on } \partial\Omega \times (0, T). \quad (1.4)$$

Published applications of scalar conservation laws that explicitly use zero-flux boundary conditions include, for example, the sedimentation of suspensions in closed vessels [3, 4, 2] and the dispersal of a single species of animals in a finite territory [18]. However, the boundary condition (1.4) is physically reasonable also in other applications, for example when (1.1) appears as the vanishing viscosity limit of a multi-dimensional model of turbulence [5] or of a simple model of two-phase flow in porous media [21].

To put the paper in the proper perspective, let us first recall some previous treatments of the Dirichlet problem (1.1), (1.2), (1.3). One major difficulty associated with this problem is due to the well-known propagation of solution values of (1.1) along characteristics, which may intersect

Date: October 29, 2004.

^aInstitut für Angewandte Analysis und Numerische Simulation, Universität Stuttgart, Pfaffenwaldring 57, D-70569 Stuttgart, Germany. E-mail: buerger@mathematik.uni-stuttgart.de.

^bInstituto de Matemática Pura e Aplicada (IMPA), Estrada Dona Castorina 110, Jardim Botânico, CEP 22460-320, Rio de Janeiro, RJ, Brazil. E-mail: hermano@impa.br.

^cCentre of Mathematics for Applications (CMA), Department of Mathematics, University of Oslo, P.O. Box 1053, Blindern, N-0316 Oslo, Norway. E-mail: kennethk@math.uio.no.

$\partial\Omega$ from the interior of Ω , such that (1.3) does not hold in a pointwise sense for all times. The well-posedness of (1.1), (1.2), (1.3) in this situation has been recovered by the use of (for example, set-valued) entropy boundary conditions. The first existence and uniqueness analysis for BV solutions of (1.1), (1.2), (1.3) is due to Bardos et al. [1]. The BV property, which is established in [1] by deriving uniform BV estimates for the solutions of a regularized (uniformly parabolic) problem, ensures the existence of boundary traces, which is crucial for the uniqueness result. It was only later that Otto [15, 19, 20] was able to study the same problem in the less restrictive L^∞ setting, for which boundary traces do not exist in general, a fact that complicates significantly the notion of solution and the proofs. See also Chen and Frid [7, 8, 9] for formulations of boundary conditions in terms of divergence-measure fields. Finally, we mention that the recent results in the L^∞ setting were extended to strongly degenerate parabolic equations by Carrillo [6], Mascia et al. [16], and Michel and Vovelle [17], while the BV approach of [1] had been transferred to this type of equations much earlier [25]. Recently Vasseur [24] showed that L^∞ entropy solutions to (1.1) always have traces at the boundaries of Q_T . This result holds for genuinely nonlinear fluxes $\mathbf{f}(u)$ (in the sense of [14]), on domains Q_T whose boundaries satisfy a mild regularity assumption, and is independent of the initial and boundary conditions. Consequently, the L^∞ case for genuinely nonlinear fluxes can be treated as in Bardos et al. [1], i.e., the more complicated notion of entropy solution used by Otto can be avoided.

Karlsen, Lie and Risebro [11] showed that a front tracking method [10] converges to a weak solution of (1.1), (1.2), (1.4) if this problem is studied in one spatial dimension. This weak solution is unique in the class of functions that can be constructed as the L^1 limit of front tracking approximations. Moreover, they present numerical results for the case of two spatial dimensions. However, for none of these cases they present a notion of entropy solution for which existence *and* uniqueness is proved.

In this paper, we suggest a notion of L^∞ entropy solutions to the zero-flux problem (1.1), (1.2), (1.4) and prove its well-posedness (existence and uniqueness) in arbitrary space dimensions. Our notion of entropy solution involves a certain boundary term in the entropy integral inequality. In fact, we can show that this entropy formulation implies that the zero-flux boundary condition is satisfied in an almost everywhere sense. The new results are valid if the flux vector satisfies the genuine nonlinearity condition of [14]. This condition is imposed to ensure the existence of boundary traces via the result of Vasseur [24]. Vasseur's result is used herein as a main tool for establishing the equivalence of two alternative definitions of entropy solutions. One of them (Definition 3) consists of the above-mentioned entropy integral inequality that incorporates the boundary term, while the other (Definition 4) states the entropy inequality in the interior of the domain and the initial and boundary conditions as separate ingredients. We mention that the fluxes used in [11] satisfy the genuine nonlinearity condition used of [14].

Let us remark that it is unclear whether the BV approach may be applied at all to the zero-flux problem. In particular, the estimating techniques of Bardos et al. [1] cannot be applied here. The difficulty is that the regularized zero-flux boundary condition does not permit control over the tangential derivatives (with respect to $\partial\Omega$) of the solution. Thus, boundary traces of solutions to (1.1), (1.2), (1.4) seem hard to obtain via BV estimates, and this has motivated the approach taken in the present paper.

The remainder of this paper is organized as follows. In Section 2 we state some technical assumptions, introduce the concepts of domains with Lipschitz deformable boundaries and traces, and recall Vasseur's result from [24]. In Section 3 we present two alternative definitions of entropy solutions to (1.1), (1.2), (1.4), and prove their equivalence by using Vasseur's result. In particular, it turns out that these entropy solutions are characterized by pointwise satisfaction of the boundary condition (1.4), in contrast to what is known for the Dirichlet problem. In Sections 4 and 5 we prove the existence and uniqueness of entropy solutions, respectively. Finally, in Section 6 we study the asymptotic behavior (for $t \rightarrow \infty$) of the entropy solutions under some additional assumptions on Ω and $\mathbf{f}(u)$.

2. ASSUMPTIONS AND PRELIMINARIES

We ask that the flux vector $\mathbf{f}(u)$ depends smoothly on u for $u \in [0, u_{\max}]$, for some fixed $u_{\max} > 0$. To ensure an L^∞ bound on the solutions, we assume that

$$\mathbf{f}(0) = 0, \quad \mathbf{f}(u_{\max}) = 0. \quad (2.1)$$

To ensure the existence of boundary traces, we assume that the flux $\mathbf{f}(u)$ is genuinely nonlinear in the following sense [14]:

$$\forall (\tau, \zeta) \in \mathbb{R} \times \mathbb{R}^N, \quad \tau^2 + |\zeta|^2 = 1 : \quad \mathcal{L}(\{u \in [0, u_{\max}] \mid \tau + \zeta \cdot \mathbf{f}'(u) = 0\}) = 0, \quad (2.2)$$

where \mathcal{L} denotes the one-dimensional Lebesgue measure. This condition is satisfied if (see [9])

$$\mathcal{L}(\{u \in [0, u_{\max}] \mid \zeta \cdot \mathbf{f}''(u) = 0\}) = 0 \quad \text{for all } \zeta \in \mathbb{R}^N \text{ with } |\zeta| = 1.$$

We adopt the usual entropy criterion, namely we consider only those weak solutions u that satisfy the inequality

$$\partial_t \eta(u) + \nabla_{\mathbf{x}} \cdot \mathbf{q}(u) \leq 0 \quad \text{on } Q_T \text{ in the sense of distributions,}$$

for every entropy pair (η, \mathbf{q}) consisting of a convex entropy function $\eta = \eta(u)$ and a corresponding entropy flux defined by $\mathbf{q}'(u) = \mathbf{f}'(u)\eta'(u)$. It is sufficient to consider the Kruřkov entropy functions $|u - k|$ along with the associated entropy fluxes $\text{sgn}(u - k)(\mathbf{f}(u) - \mathbf{f}(k))$, $k \in \mathbb{R}$.

To state Vasseur's result [24], we introduce the concept of sets with Lipschitz deformable boundaries [7]. To this end, consider an open subset $\Omega \subseteq \mathbb{R}^N$ with a boundary $\partial\Omega$.

Definition 1. *We say that $\partial\Omega$ is a deformable Lipschitz boundary provided that the following hold:*

- (a) *For all $\mathbf{x} \in \partial\Omega$ there exists a number $r > 0$ and a Lipschitz map $h : \mathbb{R}^{N-1} \rightarrow \mathbb{R}$ such that, after rotating and relabeling coordinates if necessary,*

$$\Omega \cap \mathcal{Q}(\mathbf{x}, r) = \{\mathbf{y} \in \mathbb{R}^N : h(y_1, \dots, y_{N-1}) < y_N\} \cap \mathcal{Q}(\mathbf{x}, r),$$

where $\mathcal{Q}(\mathbf{x}, r) := \{\mathbf{y} \in \mathbb{R}^N : |x_i - y_i| \leq r, i = 1, \dots, N\}$. We denote by \tilde{h} the map $(y_1, \dots, y_{N-1}) \mapsto (\tilde{\mathbf{y}}, h(\tilde{\mathbf{y}}))$.

- (b) *There exists a mapping $\Psi : \partial\Omega \times [0, 1] \rightarrow \overline{\Omega}$ such that Ψ is a homeomorphism that bi-Lipschitz over its image with $\Psi(\boldsymbol{\omega}, 0) = \boldsymbol{\omega}$ for all $\boldsymbol{\omega} \in \partial\Omega$. The map Ψ is called a Lipschitz deformation of the boundary $\partial\Omega$. We denote $\Psi_s(\boldsymbol{\omega}) = \Psi(\boldsymbol{\omega}, s)$ and $\partial\Omega_s = \Psi_s(\partial\Omega)$. We also denote by Ω_s the bounded open set whose boundary is $\partial\Omega_s$.*

Moreover, the Lipschitz deformation is said to be regular if

$$\lim_{s \rightarrow 0^+} \nabla \Psi_s \circ \tilde{h} = \nabla \tilde{h} \quad \text{in } L^1_{\text{loc}}(B), \quad (2.3)$$

where B denotes the greatest open set such that $\tilde{h}(B) \subseteq \partial\Omega$.

Obviously, if $\Omega \subseteq \mathbb{R}^N$ is an open set with a deformable Lipschitz boundary, then $Q_T = \Omega \times (0, T)$ is also an open set with deformable Lipschitz boundary in \mathbb{R}^{N+1} .

Our concept of trace is stated in the following definition.

Definition 2. *Let $Q \subseteq \mathbb{R}^{N+1}$ have a regular deformable Lipschitz boundary. We say that a given function $u \in L^\infty(Q)$ possesses a strong trace u^τ at ∂Q if $u^\tau \in L^\infty(\partial Q)$ has the property that for every regular (with respect to ∂Q) Lipschitz deformation ψ and every compact set $K \subseteq \partial Q$,*

$$\text{ess lim}_{s \rightarrow 0} \int_K |u(\psi(s, \mathbf{x})) - u^\tau(\mathbf{x})| d\mathcal{H}^N(\mathbf{x}) = 0, \quad (2.4)$$

where \mathcal{H}^N is the N -dimensional Hausdorff measure.

The following result is proved in [24].

Theorem 1. *Let $Q \subseteq \mathbb{R}^{N+1}$ have a regular deformable Lipschitz boundary, and assume that $\mathbf{f}(u)$ satisfies the genuine nonlinearity condition (2.2). Then for every function $u \in L^\infty$ satisfying the conservation law $\partial_t u + \nabla_{\mathbf{x}} \cdot \mathbf{f}(u) = 0$ in Q and the entropy inequality $\partial_t \eta(u) + \nabla_{\mathbf{x}} \cdot \mathbf{q}(u) \leq 0$ in Q for every entropy pair (η, \mathbf{q}) , the trace $u^\tau \in L^\infty(\partial Q)$ exists. In particular, $(G(u))^\tau = G(u^\tau)$ for every smooth function G .*

3. DEFINITION OF ENTROPY SOLUTIONS

From now on in this paper, it is always understood that $\Omega \subseteq \mathbb{R}^N$ in (1.1) is a bounded open set with a deformable Lipschitz boundary. Moreover, for each $T > 0$, we shall use the notation

$$Q_T := \Omega \times (0, T).$$

We denote by $C_0^\infty(Q_T)$ the set of all infinitely smooth functions vanishing on the boundary of Q_T , whereas if these functions have compact support, we write $C_c^\infty(Q_T)$ instead.

Definition 3. *A function $u \in L^\infty(Q_T)$ is called an entropy solution of the initial-boundary value problem (1.1), (1.2), (1.4) if the following entropy inequality holds:*

$$\begin{aligned} & \forall k \in \mathbb{R}, \forall \varphi \in C^\infty(\mathbb{R}^{N+1}) \text{ with } \varphi \geq 0 : \\ & \int_0^T \int_\Omega \left\{ |u - k| \partial_t \varphi + \operatorname{sgn}(u - k) (\mathbf{f}(u) - \mathbf{f}(k)) \cdot \nabla \varphi \right\} d\mathbf{x} dt \\ & + \int_\Omega \left\{ |u_0(\mathbf{x}) - k| \varphi(\mathbf{x}, 0) - |u(\mathbf{x}, T) - k| \varphi(\mathbf{x}, T) \right\} dx \\ & + \int_0^T \int_{\partial\Omega} \operatorname{sgn}(u^\tau - k) \mathbf{f}(k) \cdot \mathbf{n} \varphi(\mathbf{x}, t) d\mathcal{H}^{N-1} dt \geq 0. \end{aligned} \quad (3.1)$$

The following definition presents an alternative solution concept.

Definition 4. *A function $u \in L^\infty(Q_T)$ is called an entropy solution of the initial-boundary value problem (1.1), (1.2), (1.4) if the following conditions are satisfied:*

- (1) *The following entropy inequality is satisfied:*

$$\begin{aligned} & \forall k \in \mathbb{R}, \forall \varphi \in C_0^\infty(Q_T), \varphi \geq 0 : \\ & \int_0^T \int_\Omega \left\{ |u - k| \partial_t \varphi + \operatorname{sgn}(u - k) (\mathbf{f}(u) - \mathbf{f}(k)) \cdot \nabla \varphi \right\} d\mathbf{x} dt \geq 0. \end{aligned} \quad (3.2)$$

- (2) *The initial condition is satisfied as a limit in the following L^1 sense:*

$$\operatorname{ess\,lim}_{t \rightarrow 0^+} \int_\Omega |u(\mathbf{x}, t) - u_0(\mathbf{x})| d\mathbf{x} = 0. \quad (3.3)$$

- (3) *The boundary condition (1.4) is satisfied in the following pointwise sense:*

$$\mathbf{f}(u^\tau(\mathbf{x}, t)) \cdot \mathbf{n} = 0 \quad \text{a.e. on } \partial\Omega \times (0, T), \quad (3.4)$$

where u^τ is the trace of u , which exists thanks to Theorem 1.

Before we show that both definitions are equivalent, as is stated in Lemma 1 below, let us mention that Definition 3 will be used for the existence proof, while both Definition 3 and Definition 4 will be used for proving uniqueness.

Lemma 1. *A function $u \in L^\infty(Q_T)$ is an entropy solution in the sense of Definition 3 if and only if it is an entropy solution in the sense of Definition 4.*

Proof. We first prove that Definition 3 implies Definition 4. It is obvious that (3.1) implies (3.2). To show that (3.3) is satisfied, we choose in (3.1) the test function $\varphi(\mathbf{x}, t) = \zeta(t)\xi(\mathbf{x})$, where $\zeta \in C_c^\infty(-\infty, \delta)$, $\delta > 0$, $\xi \in C_0^\infty(\Omega)$, $\zeta \geq 0$, $\xi \geq 0$, which implies

$$\forall k \in \mathbb{R} : \int_0^\delta \zeta'(t) \int_\Omega |u - k| \xi(\mathbf{x}) d\mathbf{x} dt + \int_\Omega |u_0(\mathbf{x}) - k| \xi(\mathbf{x}) d\mathbf{x} + C \int_0^\delta \zeta(t) dt \geq 0.$$

Choosing $\zeta(t) = \chi_{(-\delta, \delta)}(t)$ (after mollifying and passing to the limit), we get for $\delta \rightarrow 0$

$$\forall k \in \mathbb{R} : \quad -\operatorname{ess\,lim}_{t \rightarrow 0^+} \int_{\Omega} |u - k| \xi(\mathbf{x}) \, d\mathbf{x} + \int_{\Omega} |u_0(\mathbf{x}) - k| \xi(\mathbf{x}) \, d\mathbf{x} \geq 0. \quad (3.5)$$

The limit on the left-hand side exists due to Theorem 1. The initial condition (3.3) follows by taking $k < 0$ and $k > u_{\max}$ in (3.5). Inequality (3.1) implies

$$\forall \varphi \in C_0^\infty(Q_T) : \quad \int_0^T \int_{\Omega} \{u \partial_t \varphi + \mathbf{f}(u) \cdot \nabla \varphi\} \, d\mathbf{x} \, dt = 0. \quad (3.6)$$

Indeed, it suffices to take $k = u_{\max}$ and $k = 0$ in (3.1), where we recall (2.1). Now we use in (3.6) the test function $\varphi(x, t) = \Phi(t) \xi(\mathbf{x}) (1 - \mu_h(\mathbf{x}))$, where $\Phi \in C_0^\infty(0, T)$, $\xi \in C_0^\infty(\bar{\Omega})$, and $\{\mu_h\}_{h>0}$ is a sequence of functions in $C^2(\Omega) \cap C(\bar{\Omega})$ such that

$$\lim_{h \rightarrow 0} \mu_h = 1 \text{ pointwise in } \Omega, \quad 0 \leq \mu_h \leq 1, \quad \mu_h = 0 \text{ on } \partial\Omega. \quad (3.7)$$

Taking the limit $h \rightarrow 0$ in the equation

$$\int_0^T \int_{\Omega} \left\{ u \Phi'(t) \xi(\mathbf{x}) (1 - \mu_h(\mathbf{x})) + \Phi(t) (1 - \mu_h(\mathbf{x})) \mathbf{f}(u) \cdot \nabla \xi(\mathbf{x}) - \Phi(t) \xi(\mathbf{x}) \mathbf{f}(u) \cdot \nabla \mu_h \right\} \, d\mathbf{x} \, dt = 0,$$

and using that the function $\xi(\mathbf{x})$ may be chosen arbitrarily, we obtain (3.4).

As for the converse, let

$$\chi^h(t) := \begin{cases} t/h & \text{for } 0 \leq t \leq h, \\ 1 & \text{for } h < t \leq T - h, \\ (T - t)/h & \text{for } T - h < t < T, \\ 0 & \text{for } t \notin (0, T). \end{cases}$$

Also, for $s \in [0, 1]$ let the function $\zeta_s \in \operatorname{Lip}(\mathbb{R}^N)$ be defined by

$$\zeta_s(\mathbf{x}) := \begin{cases} 1 & \text{for } \mathbf{x} \in \Omega_s, \\ r/s & \text{for } \mathbf{x} \in \partial\Omega_r, 0 \leq r \leq s, \\ 0 & \text{for } \mathbf{x} \notin \Omega, \end{cases}$$

where $\partial\Omega_s$ is the image of $\partial\Omega$ under the Lipschitz deformation $\Psi(\boldsymbol{\omega}, s)$ with $\partial\Omega_0 = \partial\Omega$, and Ω_s is the bounded open set whose boundary is $\partial\Omega_s$. For notational convenience, we also introduce the function $\mathbf{F}(u, k) := \operatorname{sgn}(u - k)(\mathbf{f}(u) - \mathbf{f}(k))$.

Now let us define the function $\varphi(\mathbf{x}, t) = \chi^h(t) \zeta_s(\mathbf{x}) \tilde{\varphi}(\mathbf{x}, t)$, where $\tilde{\varphi} \in C^\infty(\mathbb{R}^{N+1})$. An approximation argument reveals that we may use φ as a test function for (3.2). Then we obtain

$$\begin{aligned} \forall k \in \mathbb{R} : \quad & \int_0^T \int_{\Omega} \{ |u - k| \partial_t \tilde{\varphi} + \mathbf{F}(u, k) \cdot \nabla \tilde{\varphi} \} \, d\mathbf{x} \, dt \\ & - \int_0^T \int_{\Omega} \{ |u - k| \partial_t \tilde{\varphi} + \mathbf{F}(u, k) \cdot \nabla \tilde{\varphi} \} (1 - \chi^h(t) \zeta_s(\mathbf{x})) \, d\mathbf{x} \, dt \\ & + \int_0^T \int_{\Omega} |u - k| \zeta_s(\mathbf{x}) (\chi^h)'(t) \tilde{\varphi} \, d\mathbf{x} \, dt + \int_0^T \int_{\Omega} \chi^h(t) \tilde{\varphi} \mathbf{F}(u, k) \cdot \nabla \zeta_s(\mathbf{x}) \, d\mathbf{x} \, dt \geq 0. \end{aligned} \quad (3.8)$$

Letting $h \rightarrow 0$ and using (3.3), we get

$$\begin{aligned} \forall k \in \mathbb{R} : \quad & \int_0^T \int_{\Omega} \{ |u - k| \partial_t \tilde{\varphi} + \mathbf{F}(u, k) \cdot \nabla \tilde{\varphi} \} \, d\mathbf{x} \, dt \\ & - \int_0^T \int_{\Omega} \{ |u - k| \partial_t \tilde{\varphi} + \mathbf{F}(u, k) \cdot \nabla \tilde{\varphi} \} (1 - \zeta_s(\mathbf{x})) \, d\mathbf{x} \, dt \\ & + \int_{\Omega} \left\{ |u_0(\mathbf{x}) - k| \tilde{\varphi}(\mathbf{x}, 0) - |u(\mathbf{x}, T) - k| \tilde{\varphi}(\mathbf{x}, T) \right\} \zeta_s(\mathbf{x}) \, d\mathbf{x} \\ & + \int_0^T \int_{\Omega} \tilde{\varphi} \mathbf{F}(u, k) \cdot \nabla \zeta_s(\mathbf{x}) \, d\mathbf{x} \, dt \geq 0. \end{aligned}$$

Finally, sending $s \rightarrow 0$, using (3.4) and replacing $\tilde{\varphi}$ by the symbol φ again, we get

$$\begin{aligned} \forall k \in \mathbb{R} : \quad & \forall \varphi \in C^\infty(\mathbb{R}^{N+1}), \varphi \geq 0 : \\ & \int_0^T \int_\Omega \{ |u - k| \partial_t \varphi + \mathbf{F}(u, k) \cdot \nabla \varphi \} d\mathbf{x} dt \\ & + \int_\Omega \{ |u_0(\mathbf{x}) - k| \varphi(\mathbf{x}, 0) - |u(\mathbf{x}, T) - k| \varphi(\mathbf{x}, T) \} d\mathbf{x} \\ & + \int_0^T \int_{\partial\Omega} \operatorname{sgn}(u - k) \mathbf{f}(k) \cdot \mathbf{n} \varphi(x, t) d\mathcal{H}^{N-1} dt \geq 0, \end{aligned}$$

which is exactly (3.1). \square

4. EXISTENCE OF ENTROPY SOLUTIONS

To show the existence of an entropy solution (in the sense of the previous section), we consider as in [2] the following regularized parabolic problem for each $\varepsilon > 0$:

$$\partial_t u^\varepsilon + \nabla_{\mathbf{x}} \cdot \mathbf{f}(u^\varepsilon) = \varepsilon \Delta u^\varepsilon, \quad (\mathbf{x}, t) \in Q_T, \quad (4.1a)$$

$$u^\varepsilon(\mathbf{x}, 0) = u_0^\varepsilon(\mathbf{x}), \quad \mathbf{x} \in \Omega, \quad (4.1b)$$

$$(\mathbf{f}(u^\varepsilon) - \varepsilon \nabla_{\mathbf{x}} u^\varepsilon) \cdot \mathbf{n} = 0 \quad \text{on } \partial\Omega, \quad (4.1c)$$

where u_0^ε is a sequence of smooth functions that converges to u_0 in $L^p(\Omega)$ for $1 \leq p < \infty$ and respects the minimum/maximum values of u_0 . The existence and uniqueness of a classical solution to (4.1) follows from standard arguments, see e.g. [13, Ch. V].

Lemma 2. *Suppose (2.1) holds. Then $u_0(x) \in [0, u_{\max}]$ for a.e. $x \in \Omega$ implies $u^\varepsilon(x, t) \in [0, u_{\max}]$ for every $(x, t) \in Q_T$.*

Proof. The maximum principle shows that any extremum u^ε assumes in the interior of Q_T belongs to $[0, u_{\max}]$. Moreover, u^ε is also bounded on $\Omega \times \{0\}$ due to the assumption on u_0 . Condition (4.1c) can also be rewritten as

$$\mathbf{f}(u^\varepsilon) \cdot \mathbf{n} = \varepsilon \nabla_{\mathbf{x}} u^\varepsilon \cdot \mathbf{n} = \varepsilon \partial_{\mathbf{n}} u^\varepsilon \quad \text{on } \partial\Omega, \quad t \in (0, T], \quad (4.2)$$

where $\partial_{\mathbf{n}}$ denotes the outward normal derivative on $\partial\Omega$. Assume now that u^ε takes a local extremum at $(\mathbf{x}_0, t_0) \in \partial\Omega \times (0, T]$. To see that $u^\varepsilon(\mathbf{x}_0, t_0) \in [0, u_{\max}]$, we assume that $u^\varepsilon(\mathbf{x}_0, t_0) = M \notin [0, u_{\max}]$. Without loss of generality, let M be the maximum of u^ε on $\overline{Q_T}$. Since u^ε is smooth, we can assume that there exists a number $\rho > 0$ such that $u^\varepsilon > u_{\max}$ on $\mathcal{D} := \{\mathbf{x} \in \mathbb{R}^N : \|\mathbf{x} - \mathbf{x}_0\| < \rho\} \cap \Omega$. Consequently, u^ε satisfies the linear parabolic equation $\mathcal{L}u^\varepsilon \equiv \varepsilon \Delta u^\varepsilon - \partial_t u^\varepsilon = 0$ on \mathcal{D} . By closely following standard treatments for linear parabolic PDEs (see e.g. in [22, 23]), one can easily show that $\partial_{\mathbf{n}} u^\varepsilon(\mathbf{x}_0, t_0) > 0$ without invoking the boundary condition (4.1c). However, for $u^\varepsilon(\mathbf{x}_0, t_0) \notin [0, u_{\max}]$ we have $\mathbf{f} = 0$ and hence $\mathbf{f} \cdot \mathbf{n} = 0$. By (4.2) this implies $\varepsilon \partial_{\mathbf{n}} u^\varepsilon = 0$, i.e. $\partial_{\mathbf{n}} u^\varepsilon = 0$, a contradiction. Thus, if u^ε assumes its maximum on $\partial\Omega \times (0, T]$, then this maximum must belong to $[0, u_{\max}]$. The same argument also applies to the minimum. \square

Theorem 2. *Suppose $u_0(x) \in [0, u_{\max}]$ for a.e. $x \in \Omega$ and that conditions (2.1), (2.2) hold. Then there exists an entropy solution u of the zero-flux initial-boundary value problem (1.1), (1.2), (1.4), which moreover satisfies $u(x, t) \in [0, u_{\max}]$ for a.e. $(x, t) \in Q_T$.*

Proof. Let η_δ be a smooth convex function with a smoothing parameter $\delta > 0$, and define the corresponding entropy flux \mathbf{q}_δ by $(\mathbf{q}_\delta)' = \eta_\delta'(\mathbf{f})'$. Then let $\varphi \in C^\infty(\mathbb{R}^{N+1})$ be a nonnegative test function. Multiplying (4.1a) by $\eta_\delta'(u^\varepsilon)\varphi(\mathbf{x}, t)$, integrating the result over Q_T , using integration by parts, and using the initial condition (4.1b) yields

$$\begin{aligned} 0 &= \iint_{Q_T} \left\{ \partial_t \eta_\delta(u^\varepsilon) \varphi + \nabla \cdot \mathbf{q}(u^\varepsilon) \varphi - \varepsilon \Delta \eta_\delta(u^\varepsilon) \varphi \right\} dt d\mathbf{x} \\ &= - \int_\Omega \left\{ \eta_\delta(u_0^\varepsilon(\mathbf{x})) \varphi(\mathbf{x}, 0) - \eta_\delta(u^\varepsilon(\mathbf{x}, T)) \varphi(\mathbf{x}, T) \right\} d\mathbf{x} \end{aligned}$$

$$\begin{aligned}
 & + \int_0^T \int_{\partial\Omega} (q_\delta(u^\varepsilon) - \varepsilon \eta'_\delta(u^\varepsilon) \nabla u^\varepsilon) \cdot \mathbf{n} \varphi(\mathbf{x}, t) d\mathcal{H}^{N-1} dt \\
 & - \iint_{Q_T} \{ \eta_\delta(u^\varepsilon) \partial_t \varphi + \mathbf{q}_\delta(u^\varepsilon) \cdot \nabla \varphi + \varepsilon \nabla \eta_\delta(u^\varepsilon) \cdot \nabla \varphi \} dt d\mathbf{x} + \varepsilon \iint_{Q_T} \eta''_\delta(u^\varepsilon) |\nabla u^\varepsilon|^2 \varphi(\mathbf{x}, t) dt d\mathbf{x}.
 \end{aligned}$$

Considering that the last integral is positive and taking into account that due to (4.1c),

$$(\eta'_\delta(u^\varepsilon) \mathbf{f}(u^\varepsilon) - \varepsilon \nabla \eta_\delta(u^\varepsilon)) \cdot \mathbf{n} = 0 \quad \text{on } \partial\Omega \times (0, T),$$

we obtain the inequality

$$\begin{aligned}
 & \iint_{Q_T} \{ \eta_\delta(u^\varepsilon) \partial_t \varphi + \mathbf{q}_\delta(u^\varepsilon) \cdot \nabla \varphi \} dt d\mathbf{x} + \int_\Omega \{ \eta_\delta(u^\varepsilon_0(\mathbf{x})) \varphi(\mathbf{x}, 0) - \eta_\delta(u^\varepsilon(\mathbf{x}, T)) \varphi(\mathbf{x}, T) \} d\mathbf{x} \\
 & - \int_0^T \int_{\partial\Omega} (\mathbf{q}_\delta(u^\varepsilon) - \eta'_\delta(u^\varepsilon) \mathbf{f}(u^\varepsilon)) \cdot \mathbf{n} \varphi(\mathbf{x}, t) d\mathcal{H}^{N-1} dt \\
 & + \varepsilon \left\{ \int_0^T \int_{\partial\Omega} \eta_\delta(u^\varepsilon) \nabla \varphi \cdot \mathbf{n} d\mathcal{H}^{N-1} dt - \iint_{Q_T} \eta_\delta(u^\varepsilon) \Delta \varphi d\mathbf{x} dt \right\} \geq 0.
 \end{aligned} \tag{4.3}$$

We now first keep δ and φ fixed and take the limit $\varepsilon \rightarrow 0$. Since u^ε is bounded, the terms in the last line of (4.3) vanish. Using the compactness theorem in [14], we know that the sequence $u^\varepsilon(\mathbf{x}, t)$ is compact in $L^1_{\text{loc}}(Q_T)$. Then, making $\varepsilon \rightarrow 0$ in (4.3), passing to a subsequence if necessary, we obtain

$$\begin{aligned}
 & \iint_{Q_T} \{ \eta_\delta(u) \partial_t \varphi + \mathbf{q}_\delta(u) \cdot \nabla \varphi \} dt d\mathbf{x} + \int_\Omega \{ \eta_\delta(u_0(\mathbf{x})) \varphi(\mathbf{x}, 0) - \eta_\delta(u(\mathbf{x}, T)) \varphi(\mathbf{x}, T) \} d\mathbf{x} \\
 & - \int_0^T \int_{\partial\Omega} (\mathbf{q}_\delta(u^\tau) - \eta'_\delta(u^\tau) \mathbf{f}(u^\tau)) \cdot \mathbf{n} \varphi(\mathbf{x}, t) d\mathcal{H}^{N-1} dt \geq 0,
 \end{aligned} \tag{4.4}$$

where we denote by \mathbf{q}_δ the smoothed entropy flux defined by $\mathbf{q}'_\delta = \eta'_\delta \mathbf{f}'$.

Finally, we assume that $\eta_\delta(u)$ is a smooth approximation to $|u - k|$ as $\delta \rightarrow 0$, for example

$$\eta_\delta(z) = ((z - k)^2 + \delta^2)^{1/2} - \delta, \quad k \in \mathbb{R}, \tag{4.5}$$

so that

$$\mathbf{q}_\delta(u) - \eta'_\delta(u) \mathbf{f}(u) \xrightarrow{\delta \rightarrow 0} \mathbf{F}(u, k) - \text{sgn}(u - k) \mathbf{f}(u) = -\text{sgn}(u - k) \mathbf{f}(k). \tag{4.6}$$

Thus, letting $\delta \rightarrow 0$ and using (4.6), we obtain (3.1), which concludes the proof. \square

5. UNIQUENESS OF ENTROPY SOLUTIONS

Theorem 3. *Suppose $u_0, v_0 \in L^\infty(\Omega)$ and that conditions (2.1), (2.2) hold. Let u and v be entropy solutions of (1.1), (1.2), (1.4) with initial conditions $u|_{t=0} = u_0$ and $v|_{t=0} = v_0$, respectively. Then for any $t > 0$*

$$\int_\Omega |u(\mathbf{x}, t) - v(\mathbf{x}, t)| d\mathbf{x} \leq \int_\Omega |u_0(\mathbf{x}) - v_0(\mathbf{x})| d\mathbf{x}. \tag{5.1}$$

In particular, there exists at most one entropy solution to the zero-flux initial-boundary value problem (1.1), (1.2), (1.4).

Proof. We consider two entropy solutions $u = u(\mathbf{x}, t)$ and $v = v(\mathbf{y}, s)$. Then the standard ‘‘doubling of the variables’’ argument [12] yields that for all nonnegative functions $\varphi = \varphi(\mathbf{x}, t, \mathbf{y}, s)$ in $C^\infty(Q_T \times Q_T)$ having the property that $\varphi(\cdot, \cdot, \mathbf{y}, s), \varphi(\mathbf{x}, t, \cdot, \cdot) \in C_c^\infty(Q_T)$ for each $(\mathbf{y}, s) \in Q_T$ and $(\mathbf{x}, t) \in Q_T$, respectively, the following inequality holds:

$$\iiint_{Q_T \times Q_T} \{ |u - v| (\partial_t \varphi + \partial_s \varphi) + \mathbf{F}(u, v) \cdot (\nabla_{\mathbf{x}} \varphi + \nabla_{\mathbf{y}} \varphi) \} ds dy dt d\mathbf{x} \geq 0. \tag{5.2}$$

We pick $\theta \in C_c^\infty(0, T)$, $\theta \geq 0$, and choose in (5.2)

$$\varphi(\mathbf{x}, t, \mathbf{y}, s) := \mu_\delta(\mathbf{x}) \mu_\eta(\mathbf{y}) \rho_{l,m}(\mathbf{x}, t, \mathbf{y}, s) \theta(t), \quad \delta, \eta > 0, l, m \in \mathbb{N},$$

where μ_δ, μ_η are sequences of the type used in the proof of Lemma 1, see in particular (3.7), and

$$\rho_{l,m}(\mathbf{x}, t, \mathbf{y}, s) := \rho_l(t-s)\rho_m(\mathbf{x}-\mathbf{y}),$$

with $\{\rho_l\}_{l \in \mathbb{N}}$ and $\{\rho_m\}_{m \in \mathbb{N}}$ being sequences of symmetric mollifiers in \mathbb{R} and \mathbb{R}^N , respectively. Setting $\partial_{t+s} := \partial_t + \partial_s$, $\nabla_{\mathbf{x}+\mathbf{y}} := \nabla_{\mathbf{x}} + \nabla_{\mathbf{y}}$, we obtain from (5.2)

$$\begin{aligned} & \iiint_{Q_T \times Q_T} |u-v| \mu_\delta \mu_\eta \rho_{l,m} \theta' ds d\mathbf{y} dt d\mathbf{x} \\ & + \iiint_{Q_T \times Q_T} \mathbf{F}(u,v) (\nabla_{\mathbf{x}} \mu_\delta) \mu_\eta \rho_{l,m} \theta ds d\mathbf{y} dt d\mathbf{x} \\ & + \iiint_{Q_T \times Q_T} \mathbf{F}(u,v) \mu_\delta (\nabla_{\mathbf{y}} \mu_\eta) \rho_{l,m} \theta ds d\mathbf{y} dt d\mathbf{x} \\ & =: I_1^{\delta,\eta,l,m} + I_2^{\delta,\eta,l,m} + I_3^{\delta,\eta,l,m} \geq 0. \end{aligned} \tag{5.3}$$

It is clear that

$$I_1^{\delta,\eta,l,m} \xrightarrow{\delta,\eta \rightarrow 0} \iiint_{Q_T \times Q_T} |u-v| \rho_{l,m} \theta' ds d\mathbf{y} dt d\mathbf{x} =: I_1^{0,0,l,m}. \tag{5.4}$$

By first taking the limits $\delta, \eta \rightarrow 0$ and then taking into account that $\mathbf{f}(u^\tau) \cdot \mathbf{n} = 0$ a.e. on $\partial\Omega \times (0, T)$, we obtain

$$\begin{aligned} & I_2^{\delta,\eta,l,m} \xrightarrow{\delta,\eta \rightarrow 0} \int_0^T \int_{\partial\Omega} \iint_{Q_T} \mathbf{F}(u^\tau, v(\mathbf{y}, s)) \cdot \mathbf{n} \rho_{l,m} \theta ds d\mathbf{y} d\mathcal{H}^{N-1} dt \\ & = - \int_0^T \int_{\partial\Omega} \iint_{Q_T} \operatorname{sgn}(u^\tau - v(\mathbf{y}, s)) \mathbf{f}(v(\mathbf{y}, s)) \cdot \mathbf{n} \rho_{l,m} \theta ds d\mathbf{y} d\mathcal{H}^{N-1} dt \\ & =: I_2^{0,0,l,m}. \end{aligned} \tag{5.5}$$

In the same way, using and $\mathbf{f}(v^\tau) \cdot \mathbf{n} = 0$ a.e. on $\partial\Omega \times (0, T)$, we obtain

$$\begin{aligned} & I_3^{\delta,\eta,l,m} \xrightarrow{\delta,\eta \rightarrow 0} = - \iint_{Q_T} \int_0^T \int_{\partial\Omega} \operatorname{sgn}(u(\mathbf{x}, t) - v^\tau) \mathbf{f}(u(\mathbf{x}, t)) \cdot \mathbf{n} \rho_{l,m} \theta d\mathcal{H}^{N-1} ds dt d\mathbf{x} \\ & =: I_3^{0,0,l,m}. \end{aligned} \tag{5.6}$$

Now setting $\varphi(\mathbf{x}, t) = \rho_{l,m}(\mathbf{x}, t, \mathbf{y}, s)\theta(t)$ in (3.1) (with \mathbf{y} and s considered as parameters), we get

$$\begin{aligned} & - \int_0^T \int_{\partial\Omega} \operatorname{sgn}(u^\tau - k) \mathbf{f}(k) \cdot \mathbf{n} \rho_{l,m} \theta d\mathcal{H}^{N-1} dt \\ & \leq \iint_{Q_T} |u-k| \rho_{l,m} \theta' dt d\mathbf{x} \\ & + \iint_{Q_T} \{ |u-k| \partial_t \rho_{l,m} \theta + \mathbf{F}(u, k) \cdot \nabla_{\mathbf{x}} \rho_{l,m} \theta \} dt d\mathbf{x}, \quad \forall k \in \mathbb{R}. \end{aligned}$$

Setting $k = v(\mathbf{y}, s)$ and integrating the result over $(\mathbf{y}, s) \in Q_T$, we obtain

$$\begin{aligned} & I_2^{0,0,l,m} \leq \iiint_{Q_T \times Q_T} \{ |u-v| \rho_{l,m} \theta' ds d\mathbf{y} dt d\mathbf{x} \\ & + \iiint_{Q_T \times Q_T} \{ |u-v| \partial_t \rho_{l,m} \theta + \mathbf{F}(u, v) \cdot \nabla_{\mathbf{x}} \rho_{l,m} \theta \} ds d\mathbf{y} dt d\mathbf{x}. \end{aligned} \tag{5.7}$$

In a similar way, using the test function $\varphi(\mathbf{y}, s) = \rho_{l,m}(\mathbf{x}, t, \mathbf{y}, s)\theta(t)$ in the analogue of (3.1) for the entropy solution $v = v(\mathbf{y}, s)$ (with \mathbf{x} and t considered as parameters) and taking into account that θ is a function of t only, we get

$$\begin{aligned} I_3^{0,0,l,m} &= - \iint_{Q_T} \int_0^T \int_{\partial\Omega} \operatorname{sgn}(v^\tau - u(\mathbf{x}, t)) \mathbf{f}(u(\mathbf{x}, t)) \cdot \mathbf{n} \rho_{l,m} \theta \, d\mathcal{H}^{N-1} \, dt \, d\mathbf{x} \\ &\leq I_4^{l,m} + \iiint_{Q_T \times Q_T} \{|u - v| \partial_s \rho_{l,m} \theta + \mathbf{F}(u, v) \cdot \nabla_{\mathbf{y}} \rho_{l,m} \theta\} \, ds \, d\mathbf{y} \, dt \, d\mathbf{x}, \end{aligned} \quad (5.8)$$

where

$$\begin{aligned} I_4^{l,m} &:= \iint_{Q_T} \int_{\Omega} \left\{ |v_0(\mathbf{y}) - u(\mathbf{x}, t)| \rho_l(t) \right. \\ &\quad \left. - |v(\mathbf{y}, T) - u(\mathbf{x}, t)| \rho_l(t - T) \right\} \rho_m(\mathbf{x} - \mathbf{y}, t) \theta(t) \, d\mathbf{y} \, dt \, d\mathbf{x}. \end{aligned}$$

Combining (5.7) and (5.8) and using that $(\partial_t + \partial_s)\rho_{l,m} = 0$ and $(\nabla_{\mathbf{x}} + \nabla_{\mathbf{y}})\rho_{l,m} = 0$, we get

$$I_2^{0,0,l,m} + I_3^{0,0,l,m} \leq I_1^{0,0,l,m} + I_4^{l,m}.$$

Thus, for $\delta, \eta \rightarrow 0$ we obtain from (5.3) the inequality

$$2I_1^{0,0,l,m} + I_4^{l,m} \geq 0. \quad (5.9)$$

Next, we pass to the limits $l, m \rightarrow \infty$. Since $\theta(0) = \theta(T) = 0$, we obtain

$$I_4^{l,m} \xrightarrow{l \rightarrow \infty} \iint_{\Omega \times \Omega} \left\{ |v_0(\mathbf{y}) - u_0(\mathbf{x})| \theta(0) - |v(\mathbf{y}, T) - u(\mathbf{y}, T)| \theta(T) \right\} \rho_m(\mathbf{x} - \mathbf{y}) \, d\mathbf{y} \, d\mathbf{x} = 0.$$

Collecting the limits, we obtain the following inequality from (5.9):

$$\iint_{Q_T} |u - v| \theta' \, dt \, d\mathbf{x} \geq 0, \quad \forall \theta \in C_c^\infty(0, T). \quad (5.10)$$

Inequality (5.1) follows now from (5.10) in a standard way. \square

6. ASYMPTOTIC BEHAVIOR OF ENTROPY SOLUTIONS

In this section, we suppose that the following additional assumptions are satisfied.

- (A1) There exists a direction, given by an unity vector $\mathbf{e} \in \mathbb{R}^N$, such that the hyperplanes $\Pi_\nu = \{\mathbf{x} \in \mathbb{R}^N : \mathbf{e} \cdot \mathbf{x} = \nu\}$ cut out Ω in open sets with Lipschitz deformable boundaries, $\Omega = \Omega^1(\nu) \cup \dots \cup \Omega^{j_\nu}(\nu)$ for $\nu_{\min} < \nu < \nu_{\max}$, and do not intersect Ω for $\nu \notin [\nu_{\min}, \nu_{\max}]$.
- (A2) $\mathbf{f}(u) \cdot \mathbf{e} > 0$, for $0 < u < u_{\max}$, and $\mathbf{f}(u_{\min}) \cdot \mathbf{e} = \mathbf{f}(u_{\max}) \cdot \mathbf{e} = 0$.

Without loss of generality, we assume that \mathbf{e} is the unitary vector in the direction of the x_1 -axis. Also, for simplicity, we assume $u_{\max} = 1$.

Before we continue, let us point out that all results obtained in the previous sections hold with $T = \infty$, in which case we use the notation Q for $\Omega \times (0, \infty)$.

Theorem 4. *Assume that (A1) and (A2) hold and let $u(\mathbf{x}, t)$ be the entropy solution of (1.1), (1.2), (1.4) on Q . Then, for any $g \in C_{\text{per}}([0, 1])$ and $h > 0$, we have*

$$\lim_{t \rightarrow \infty} \int_t^{t+h} \int_{\Omega} g(u(\mathbf{x}, s)) \, d\mathbf{x} \, ds = h |\Omega| g(0). \quad (6.1)$$

Here, $C_{\text{per}}([0, 1])$ denotes the space of the continuous periodic functions in $[0, 1]$, and $|\Omega|$ denotes the measure of Ω .

Proof. Let $\Omega(\nu)$ be the union of the sets $\Omega^j(\nu)$ which lay on the left-hand (negative) side of Π_ν . Integrating (1.1) on $\Omega(\nu) \times (0, t)$, using the Gauss-Green formula [7], and (1.4), we arrive at

$$\int_{\Omega(\nu)} u(\mathbf{x}, t) \, d\mathbf{x} - \int_{\Omega(\nu)} u_0(\mathbf{x}) \, d\mathbf{x} + \int_0^t \int_{\Pi_\nu \cap \Omega} f_1(u(\mathbf{x}, s)) \, d\mathcal{H}^{N-1} \, ds = 0. \quad (6.2)$$

Integrating (6.2) with respect to ν from ν_{\min} to ν_{\max} , recalling (A2) we obtain

$$0 < \int_0^\infty \int_\Omega f_1(u(\mathbf{x}, s)) \, d\mathbf{x} \, ds \leq C$$

for some positive constant C . In particular, we have

$$\lim_{t \rightarrow \infty} \int_t^{t+h} \int_\Omega f_1(u(\mathbf{x}, s)) \, d\mathbf{x} \, ds = 0. \quad (6.3)$$

Again recalling (A2), we immediately obtain from (6.3) that the probability measures defined by

$$\langle \mu_t, g \rangle := \frac{1}{h|\Omega|} \int_t^{t+h} \int_\Omega g(u(\mathbf{x}, s)) \, d\mathbf{x} \, ds, \quad g \in C_{\text{per}}([0, 1]), \quad (6.4)$$

satisfy $\mu_t \rightarrow \delta_0$ in the weak- \star topology of $C_{\text{per}}([0, 1])^*$, and so (6.1) follows. \square

ACKNOWLEDGMENTS

R. Bürger was supported by the Collaborative Research Programme (Sonderforschungsbereich) 404 “Mehrfeldprobleme in der Kontinuumsmechanik” at the University of Stuttgart. H. Frid was partially supported by CNPq through the grants 352871/96-2, 46.5714/00-5, 479416/2001-0, and FAPERJ through the grant E-26/151.890/2000. K. H. Karlsen was supported by the Research Council of Norway through an Outstanding Young Investigators Award and by the HYKE network through the EC as contract HPRN-CT-2002-00282.

REFERENCES

- [1] C. Bardos, A.Y. Le Roux, J.C. Nédélec, First order quasilinear equations with boundary conditions, *Comm. PDE* 4 (1979) 1017–1034.
- [2] R. Bürger, S. Evje, K.H. Karlsen, On strongly degenerate convection-diffusion problems modeling sedimentation-consolidation processes, *J. Math. Anal. Appl.* 247 (2000) 517–556.
- [3] R. Bürger, K.H. Karlsen, On some upwind schemes for the phenomenological sedimentation-consolidation model, *J. Eng. Math.* 41 (2001) 145–166.
- [4] R. Bürger, M. Kunik, A critical look at the kinematic-wave theory for sedimentation-consolidation processes in closed vessels, *Math. Meth. Appl. Sci.* 24 (2001) 1257–1273.
- [5] J.M. Burgers, *The Nonlinear Diffusion Equation*, Reidel, Dordrecht, 1974.
- [6] J. Carrillo, Entropy solutions for nonlinear degenerate problems, *Arch. Rational Mech. Anal.* 147 (1999) 269–361.
- [7] G.-Q. Chen, H. Frid, Divergence-measure fields and hyperbolic conservation laws, *Arch. Rat. Mech. Anal.* 147 (1999) 89–118.
- [8] G.-Q. Chen, H. Frid, Extended divergence-measure fields and the Euler equations for gas dynamics, *Comm. Math. Phys.* 236 (2003) 251–280.
- [9] G.-Q. Chen, H. Frid, Large-time behavior of entropy solutions in L^∞ for multidimensional conservation laws. In: *Advances in Nonlinear Partial Differential Equations and Related Areas*, A volume in honor of Professor Xiqi Ding, World Scientific, Singapore 1998, 28–44.
- [10] H. Holden, N.H. Risebro, *Front Tracking for Hyperbolic Conservation Laws*, Springer-Verlag, New York 2002.
- [11] K.H. Karlsen, K.-A. Lie, N.H. Risebro, A front tracking method for conservation laws with boundary conditions. In: M. Fey and R. Jeltsch (Eds.), *Hyperbolic Problems: Theory, Numerics, Applications*, Birkhäuser Verlag, Basel 1999, 493–502.
- [12] S.N. Kružkov, First order quasilinear equations in several independent variables, *Math. USSR Sbornik* 10 (1970) 217–243.
- [13] O.A. Ladyženskaja, V.A. Solonnikov, N.N. Ural’ceva, *Linear and Quasi-linear Equations of Parabolic Type*, American Mathematical Society, Providence, RI, 1968.
- [14] P.-L. Lions, B. Perthame, E. Tadmor, A kinetic formulation of multidimensional scalar conservation laws and related equations, *J. Amer. Math. Soc.* 7 (1994) 169–191.
- [15] J. Málek, J. Nečas, M. Rokyta, M. Ružička, *Weak and Measure-Valued Solutions to Evolutionary PDEs*, Chapman & Hall, London, 1996.
- [16] C. Mascia, A. Poretta, A. Terracina, Nonhomogeneous Dirichlet problems for degenerate parabolic-hyperbolic equations, *Arch. Rat. Mech. Anal.* 163 (2002) 87–124.
- [17] A. Michel, J. Vovelle, Entropy formulation for parabolic degenerate equations with general Dirichlet boundary conditions and application to the convergence of FV methods, *SIAM J. Numer. Anal.* 41 (2003) 2262–2293.
- [18] J.D. Murray, *Mathematical Biology. II. Spatial Models and Biomedical Applications*, 3rd ed., Springer Verlag, New York 2003.

- [19] F. Otto, First order equations with boundary conditions. Preprint 234, Sonderforschungsbereich 256, University of Bonn, Germany, 1992.
- [20] F. Otto, Initial-boundary value problem for a scalar conservation law, C. R. Acad. Sci. Paris Sér. I 322 (1996) 729–734.
- [21] D.W. Peaceman, Fundamentals of Numerical Reservoir Simulation, Elsevier, Amsterdam 1977.
- [22] M. Renardy, R.C. Rogers, An Introduction to Partial Differential Equations. Springer-Verlag, New York 1993.
- [23] J. Smoller, Shock Waves and Reaction-Diffusion Equations. Springer-Verlag, New York 1983.
- [24] A. Vasseur, Strong traces for solutions of multidimensional scalar conservation laws, Arch. Rat. Mech. Anal. 160 (2001) 181–193.
- [25] Z. Wu, J. Zhao, The first boundary value problem for quasilinear degenerate parabolic equations of second order in several space variables, Chin. Ann. of Math. 4B (1983) 57–76.

Erschienene Preprints ab Nummer 2004/001

Komplette Liste: <http://preprints.ians.uni-stuttgart.de>

- 2004/001 *Geis, W., Mishuris, G., Sändig, A.-M.*: 3D and 2D asymptotic models for piezoelectric stack actuators with thin metal inclusions
- 2004/002 *Klimke, A., Wohlmuth, B., Willner, K.*: Computing expensive multivariate functions of fuzzy numbers using sparse grids
- 2004/003 *Klimke, A., Wohlmuth, B., Willner, K.*: Uncertainty modeling using efficient fuzzy arithmetic based on sparse grids: applications to dynamic systems
- 2004/004 *Flemisch, B., Mair, M., Wohlmuth, B.*: Nonconforming discretization techniques for overlapping domain decompositions
- 2004/005 *Sändig, A.-M.*: Vorlesung Mathematik für Informatiker und Softwaretechniker I, WS 2003/2004
- 2004/006 *Bürger, R., Karlsen, K. H., Towers, J. D.*: Closed-form and finite difference solutions to a population balance model of grinding mills
- 2004/007 *Berres, S., Bürger, R., Tory, E. M.*: Applications of Polydisperse Sedimentation Models
- 2004/008 *Bürger, R., Karlsen, K. H., Towers, J. D.*: A model of continuous sedimentation of flocculated suspensions in clarifier-thickener units
- 2004/009 *Bürger, R., Karlsen, K. H., Towers, J. D.*: Mathematical model and numerical simulation of the dynamics of flocculated suspensions in clarifier-thickeners
- 2004/010 *Lehrstühle: Wendland, Wohlmuth, Abteilungen: Gekeler. Sändig.*: Jahresbericht 2003
- 2004/011 *Sändig, A.-M. (Hrsg.), Knees, D. (Hrsg.)*: Nichtlineare Funktionalanalysis mit Anwendungen in der Festkörpermechanik
- 2004/012 *Wendland, W.L.*: Vorlesungsskript Partielle Differentialgleichungen
- 2004/013 *Steinbach, O. (ed.)*: Seminarbericht: Hierarchische Matrizen
- 2004/014 *Sändig, A.-M.*: Vorlesung Mathematik für Informatiker und Softwaretechniker II, SS 2004
- 2004/015 *Langer, U., Steinbach, O., Wendland, W. L. (eds)*: Workshop on Adaptive Fast Boundary Element Methods in Industrial Applications, Söllerhaus, 29.9.-2.10.2004.
- 2004/016 *Steinbach, O.*: Vorlesung Hierarchische Matrizen
- 2004/017 *Bürger, R. (Hg.)*: Seminarbericht Einführung in die Mathematische Biologie
- 2004/018 *Bürger, R., Frid, H., Karlsen, K. H.*: On the well-posedness of entropy solutions to conservation laws with a zero-flux boundary condition